

JOURNAL OF ALGEBRA **139**, 155–158 (1991)

Primordial Subgroups for $\text{mod}(kG)$

ALBERTO RAGGI-CÁRDENAS

*Instituto de Matemáticas, UNAM, Circuito Exterior,
Ciudad Universitaria, México 04510 D.F., México*

Communicated by Walter Feit

Received August 23, 1989

0. INTRODUCTION

Recall from [5] that, given a finite group G and a ring G -functor R , a subgroup H of G is called a primordial subgroup for R if

$$\sum_{L \subseteqq H} t_L^H(R(L)) \subsetneq R(H).$$

Let $\mathcal{P}(R)$ denote the set of primordial subgroups of R and let $\overline{\mathcal{P}(R)}$ denote the subgroup closure of $\mathcal{P}(R)$. In [5, 2.1], it is proved that $\overline{\mathcal{P}(R)}$ is the minimal set \mathcal{D} of subgroups of G , closed under conjugation and formation of subgroups, satisfying the condition

$$\sum_{H \in \mathcal{D}} t_H^G(R(H)) = R(G).$$

Let k be any field and consider the ring G -functor R_k , which assigns to each $H \subseteq G$ the Grothendieck ring of the category of kH -modules with the usual restriction, induction (denoted by t) and conjugation. It is well known that for fields k of zero characteristic, $\mathcal{P}(R_k)$ equals the set of k -elementary subgroups of G ; Thévenaz conjectured in [5] that this is also the case for fields of positive characteristic (see definitions below). In this work we give a short proof of this statement. First we show for the ring G -functor ${}_{\mathbb{Q}}R_k = \mathbb{Q} \otimes_{\mathbb{Z}} R_k$ that $\mathcal{P}({}_{\mathbb{Q}}R_k)$ equals the set of cyclic subgroups of G of order prime to the characteristic of k . Then we use this to prove Thévenaz conjecture.

1. NOTATION AND PRELIMINARY LEMMAS

Fix a field k of characteristic $p \geq 0$. Let n be a positive integer (prime to p if $p > 0$) and let ω_n be an n th-primitive root of unity in some algebraic closure of k . Let $k_n = k[\omega_n]$. Then k_n/k is a Galois extension of k . Let $\mathcal{G}_n =$

$\text{Gal}(k_n/k)$. Then there is a monomorphism $\mathcal{G}_n \rightarrow (\mathbb{Z}/n\mathbb{Z})^*$. We consider \mathcal{G}_n as a subgroup of $(\mathbb{Z}/n\mathbb{Z})^*$.

We say that a finite group H is k -elementary if $H = \langle x \rangle \rtimes Q$, where Q is a q -group (q a prime number), the order m of x is relatively prime to q (and to p if $p > 0$), and for each $a \in Q$ there is $t \in \mathcal{G}_m$ such that $a^{-1}xa = x^t$ (c.f. [1, Sect. 21A]).

Note that in the last definition we can replace \mathcal{G}_m by \mathcal{G}_n , where n is any multiple of m (prime to p if $p > 0$). From now on, G denotes a fixed finite group, $p > 0$, n denotes the p' -part of $|G|$, and k_0 is the prime field of k .

The next lemma tells us that we can restrict ourselves to finite fields and therefore to p -modular systems in the sense of [1, Sect. 16A].

LEMMA 1. *Let $k' = k_0[\omega_n]$ and $E = k \cap k'$. Then $R_E \cong R_k$ as ring G -functors (the isomorphism sends the class of the EG -module M to the class of $k \otimes_E M$).*

Proof. Let $H \subseteq G$ and let M_1, \dots, M_s be a set of representatives of the isomorphism classes of the simple $k'H$ -modules. Then $k_n \otimes_{k'} M_1, \dots, k_n \otimes_{k'} M_s$ is a set of representatives of the isomorphism classes of the simple $k_n H$ -modules. Clearly $\mathcal{G}_n \cong \text{Gal}(k'/E)$. The group $\text{Gal}(k'/E)$ acts on the simple $k'H$ -modules and the orbits correspond to the simple EH -modules; in fact, if \mathcal{O} is an orbit then $N_{\mathcal{O}} := \bigoplus_{M \in \mathcal{O}} M$ is a simple EG -module (see [2, 74.9]). The kH -simple modules are obtained similarly from the $k_n H$ -modules. Therefore if N_1, \dots, N_t is a set of representatives of the isomorphism classes of the EH -modules then $k \otimes_E N_1, \dots, k \otimes_E N_t$ is a set of representatives of the isomorphism classes of the kH -modules, so the lemma is proved. ■

In order to calculate $\mathcal{P}(R_k)$ we may assume that $k \subseteq k_0[\omega_n]$. Then there is a p -modular system (K, R, k) , with R a complete discrete valuation ring and K (a finite extension of the p -adic field) the quotient ring of R . There is another p -modular system (K_n, R_n, k_n) , where $K_n = K[\omega'_n]$ and $R_n = R[\omega'_n]$, with ω'_n an n th-root of unity mapping onto ω_n . It is known that $\text{Gal}(K_n/K) \cong \mathcal{G}_n$.

LEMMA 2. *The primordial set $\mathcal{P}({}_{\mathbb{Q}}R_k)$ equals the set \mathcal{C} of cyclic subgroups of G of order prime to p .*

Proof. First assume $k = k_n$. From the Dress Induction Theorem (see [6]) and the isomorphism of ring G -functors $R_k \cong \text{Brch}(G)$ (see [1, 17.14]), where $\text{Brch}(G)$ denotes the ring of Brauer characters of G , we deduce that

$${}_{\mathbb{Q}}R_k(G) = \sum_{H \in \mathcal{C}} i_H^G({}_{\mathbb{Q}}R_k(K)).$$

Therefore, by minimality, we have $\mathcal{P}({}_{\mathbb{Q}}R_k) \subseteq \mathcal{C}$.

Now, for $H \in \mathcal{C}$, since p does not divide $|H|$, we have that R_k is isomorphic as a ring H -functor to the ordinary character ring and it is known that H itself is primordial for ${}_{\mathbb{Q}}R_k$ (see [4, 3.13]). Therefore $H \in \mathcal{P}({}_{\mathbb{Q}}R_k)$.

In general there is a monomorphism of G -functors ${}_{\mathbb{Q}}R_k \hookrightarrow {}_{\mathbb{Q}}R_{k_n}$. Therefore by [5, 5.2], $\mathcal{P}({}_{\mathbb{Q}}R_k) = \mathcal{P}({}_{\mathbb{Q}}R_{k_n}) = \mathcal{C}$. ■

2. PROOF OF THE THEOREM: $\mathcal{P}(R_k) = \mathcal{E}_k$.

Consider the decomposition map of ring G -functors $d: R_K \rightarrow R_k$ (see [1, Sect. 16]). Then by [5, 2.5], $\mathcal{P}(R_k) \subseteq \mathcal{P}(R_K)$; by [1, 21.6] and [3], the latter equals \mathcal{E}_K . On the other hand for all $H \subseteq G$ it is easy to prove that the ring ${}_{\mathbb{Q}}R_k(H)$ has no nilpotent elements (see [1, Sect. 17, Exercise 4]) and so the inclusion of ring G -functors $R_k \hookrightarrow {}_{\mathbb{Q}}R_k$ together with Lemma 2 give $\mathcal{P}(R_k) \subseteq \mathcal{H}(\mathcal{C})$, where $\mathcal{H}(\mathcal{C})$ is the set of hyperelementary subgroups of G whose normal cyclic subgroup is in \mathcal{C} (see [5, 5.1]).

We claim that $\mathcal{H}(\mathcal{C}) \cap \mathcal{E}_K = \mathcal{E}_k$. The inclusion $\mathcal{H}(\mathcal{C}) \cap \mathcal{E}_K \supseteq \mathcal{E}_k$ is trivial. Suppose $H \in \mathcal{H}(\mathcal{C}) \cap \mathcal{E}_K$. Then, since $H \in \mathcal{H}(\mathcal{C})$, we have $H = \langle x \rangle \rtimes Q$, where Q is a q -group and p and q do not divide the order of x . On the other hand, since $H \in \mathcal{E}_K$, then $H = \langle y \rangle \rtimes D$, where D is a p_0 group and p_0 (a prime number) does not divide the order of y . We have three cases:

(a) $p_0 = p$. In this case p does not divide the order of y and so $H \in \mathcal{E}_k$ by the remarks above.

(b) $p_0 = q$. In this case the order of x equals the order of y , so p does not divide the order of y and so, as in (a), $H \in \mathcal{E}_k$.

(c) $p_0 \neq p$ and $p_0 \neq q$. In this case there exists $g \in H$ such that ${}^gD \subseteq \langle x \rangle$ and, since clearly gD is a characteristic subgroup of $\langle x \rangle$ which is normal in H , we conclude that D itself is normal in H , so H is cyclic and therefore $H \in \mathcal{E}_k$. This proves the claim.

Now let $H \in \mathcal{E}_k$, say, $H = \langle x \rangle \rtimes Q$ with Q a q -group. If $q \neq p$ then the decomposition map $d: R_K \rightarrow R_k$ is an isomorphism of ring H -functors. Therefore we have an isomorphism

$$\frac{R_K(H)}{\sum_{L \subsetneq H} t_L^H(R_K(L))} \cong \frac{R_k(H)}{\sum_{L \subsetneq H} t_L^H(R_k(L))}.$$

Since it is known that the left-hand side is different from zero, we have $H \in \mathcal{E}_k$.

Now assume $q = p$ and let $L \subsetneq H$. Let us prove that for each $\varphi \in R_k(L)$ we have $t_L^H(\varphi)(x) \in p\mathbb{Z}[\omega_n]$. (Here we consider the elements of $R_k(L)$ as

Brauer characters.) From the transitivity of induction we can assume without loss of generality that either $L = \langle x \rangle \rtimes Q_1$ with $Q_1 \subsetneq Q$ or else $L = \langle x^r \rangle \rtimes Q$ with $\langle x^r \rangle \subsetneq \langle x \rangle$. In the first case

$$t_L^H(\varphi)(x) = \sum_{a \in Q/Q_1} \varphi(a^{-1}xa) = \varphi(x)[Q : Q_1] \in p\mathbb{Z}[\omega_n];$$

the last equality holds since φ is constant on k -conjugacy classes (see [1; step 2 in the proof of 21.25]). In the second case $t_L^H(\varphi)(x)$ is clearly zero. We conclude that $H \in \mathcal{P}(R_k)$.

REFERENCES

1. C. W. CURTIS AND I. REINER, "Methods of Representation Theory," Vol. I, Wiley, New York, 1981.
2. C. W. CURTIS AND I. REINER, "Methods of Representation Theory," Vol. II, Wiley, New York, 1987.
3. F. LORENZ, Die Umkehrung des Satzes von Brauer-Berman-Witt über induzierte Charaktere endlichen Gruppen, *Math. Z.* **142** (1975), 167-171.
4. L. PUIG, Pointed groups and construction of modules, *J. Algebra* **116** (1988), 7-129.
5. J. THÉVENAZ, Some remarks on G -functors and the Brauer morphism, *J. Reine Angew. Math.* **384** (1988), 24-56.
6. T. YOSHIDA, Idempotents of Burnside rings and Dress induction theorem, *J. Algebra* **80** (1983), 90-105.